

DEVELOPMENT OF HIGH-FIBER PASTA USING COMPOSITE FLOUR FORMULATIONS

Anaya Zahra

BS Student, Department of English Literature, Punjab University Lahore

ABSTRACT

The development of high-fiber pasta using composite flour formulations is an important approach for improving the nutritional quality of conventional pasta while maintaining acceptable sensory and technological properties. Pasta is widely consumed because of its convenience, affordability, versatility, and relatively long shelf life; however, traditional pasta is generally produced from refined or semolina-based wheat flour and may provide limited dietary fiber. Increasing consumer awareness regarding digestive health, weight management, blood glucose regulation, and the prevention of diet-related diseases has created a growing demand for fiber-enriched food products. The present study focuses on the development of high-fiber pasta by incorporating different fiber-rich composite flour ingredients into a conventional flour formulation. Composite flour technology allows the nutritional characteristics of pasta to be improved by combining wheat flour with cereal, pseudocereal, legume, or other plant-based flour sources. The study proposes the development of pasta formulations with varying levels of composite flour and evaluates their effects on nutritional, cooking, physical, and sensory characteristics. The research uses an experimental approach in which different formulations are prepared and compared with a control pasta sample. Parameters such as cooking time, cooking loss, water absorption, texture, appearance, flavor, taste, and overall acceptability are considered. The expected findings indicate that moderate incorporation of fiber-rich flour can enhance dietary fiber content while maintaining acceptable cooking and sensory properties. However, excessive incorporation may negatively affect dough development, texture, color, and consumer acceptability. The study highlights the potential of composite flour formulations for developing nutritionally improved pasta products and provides a practical basis for producing functional pasta from locally available plant-based ingredients.

KEYWORDS: *High-fiber pasta, composite flour, dietary fiber, wheat flour, functional foods, nutritional quality, pasta formulation, sensory evaluation, food product development, healthy foods.*

INTRODUCTION

Pasta is one of the most popular cereal-based foods consumed in many parts of the world because of its convenient preparation, desirable sensory characteristics, versatility, affordability, and relatively long shelf life. Traditionally, pasta is manufactured primarily from durum wheat semolina or refined wheat flour, water, and sometimes eggs or other ingredients depending on the product type. Pasta can be prepared in numerous forms and served with different sauces, vegetables, meats, legumes, and other food ingredients, making it an important component of both household and commercial diets. Although conventional pasta provides carbohydrates, protein, and several micronutrients, its nutritional profile can be improved by increasing its dietary fiber content. The development of nutritionally enhanced pasta is therefore receiving increasing attention within food science and nutrition research. Dietary fiber is an important component of a healthy diet because it contributes to normal gastrointestinal function and can influence several physiological processes. Adequate

fiber intake is associated with improved bowel regularity and may contribute to better control of blood glucose and blood lipid levels. Fiber can also increase satiety, which may be useful in dietary strategies related to weight management. Despite these benefits, many populations consume less dietary fiber than recommended. One reason is the frequent consumption of highly refined cereal products in which bran and germ fractions have been removed during processing. Consequently, food scientists have increasingly investigated methods of incorporating fiber-rich ingredients into commonly consumed foods. Pasta represents an attractive food vehicle for fiber enrichment because it is already widely accepted by consumers.

Instead of requiring consumers to adopt completely unfamiliar foods, nutritional improvement of pasta can provide additional nutrients within a familiar product. The use of composite flour is one potential strategy for achieving this objective. Composite flour refers to a mixture of two or more flours obtained from different cereal, legume, pseudocereal, vegetable, fruit, or other plant sources. By combining wheat flour with fiber-rich ingredients, researchers can modify the nutritional and functional properties of pasta while attempting to maintain acceptable product quality. THAT FOUND .The nutritional characteristics of composite flour depend strongly on the ingredients selected. Whole-grain cereals contain bran and germ fractions that can contribute dietary fiber, minerals, vitamins, and bioactive compounds. Legume flours may provide protein, dietary fiber, resistant starch, minerals, and other phytochemicals. Ingredients such as oat flour, barley flour, chickpea flour, lentil flour, millet flour, sorghum flour, and other plant-based materials have therefore attracted attention for functional food development. The choice of flour depends not only on nutritional composition but also on technological properties such as water absorption, dough formation, mixing behavior, drying characteristics, and cooking quality.

The development of high-fiber pasta presents several technological challenges. Wheat-based pasta has particular structural characteristics because gluten proteins contribute to dough formation and provide a continuous protein network. When wheat flour is partially replaced with non-wheat composite flours, the amount and quality of gluten may decrease. This can influence dough elasticity, firmness, cooking stability, and the final texture of pasta. Fiber itself can also interact with water and other components of the dough. Because dietary fiber has considerable water-holding capacity, increasing fiber levels may alter dough hydration requirements and cooking behavior. The level of composite flour incorporation is therefore an important consideration in pasta formulation. A small or moderate replacement level may increase fiber content while preserving acceptable product characteristics. In contrast, a high replacement level may produce darker color, rougher surface, reduced elasticity, increased cooking loss, or a softer and less desirable texture. Therefore, product development requires systematic evaluation of different formulations rather than simply maximizing the amount of fiber-rich flour. Consumer acceptability is another important factor in the development of functional pasta. A product may have excellent nutritional characteristics but have limited

commercial potential if consumers dislike its appearance, taste, aroma, or texture. Sensory evaluation is consequently an essential part of high-fiber pasta development. Panelists can evaluate characteristics such as color, appearance, aroma, taste, texture, mouthfeel, and overall acceptability using a structured hedonic scale. Sensory results can help identify the formulation that provides the most appropriate balance between nutritional enhancement and consumer preference.

Incorporating such ingredients into pasta can provide opportunities for food diversification and product innovation. It may also contribute to better utilization of agricultural resources and the development of affordable functional foods. High-fiber pasta may also have potential relevance to modern dietary patterns. Consumers are increasingly interested in foods that provide nutritional benefits beyond basic energy and macronutrient supply. Functional food development has therefore expanded from specialized nutritional products toward familiar everyday foods. Pasta enriched with fiber-rich composite flour can fit within this trend because it combines a conventional food format with improved nutritional characteristics. Another important aspect of this research is the relationship between fiber enrichment and cooking properties. Pasta is generally boiled before consumption, and cooking can significantly affect its texture, water absorption, cooking loss, and nutritional characteristics. High-fiber formulations may absorb more water because of the water-binding properties of fiber. Changes in cooking time and water absorption may influence consumer perception of pasta quality. Therefore, cooking characteristics must be considered alongside nutritional analysis and sensory evaluation. The development of composite flour pasta also provides opportunities to explore sustainable food systems. Plant-based ingredients can be sourced from diverse crops, reducing dependence on a single cereal raw material. The use of locally available legumes and whole grains can support agricultural diversification and potentially increase the value of underutilized crops. However, sustainability considerations must be integrated with food safety, nutritional quality, consumer acceptance, and economic feasibility.

In food product development, formulation optimization is essential. Different proportions of wheat and composite flour can produce significantly different outcomes. The control formulation provides a reference point against which modified formulations can be compared. Experimental formulations can then be evaluated for their nutritional, physicochemical, cooking, and sensory properties. Such systematic comparison helps determine the level of substitution that produces a nutritionally improved pasta without unacceptable deterioration in quality. The present study therefore focuses on the development of high-fiber pasta using composite flour formulations. The study aims to investigate how the incorporation of fiber-rich composite flour affects the characteristics of pasta and to identify a formulation that provides improved nutritional value while retaining desirable product quality. The research combines food formulation principles, nutritional considerations, cooking evaluation, and sensory assessment to provide a comprehensive understanding of

high-fiber pasta development. Overall, the development of high-fiber pasta represents a promising strategy for improving the nutritional quality of a familiar cereal-based food. Through appropriate selection and proportioning of composite flour ingredients, it may be possible to increase dietary fiber intake without substantially compromising consumer acceptability. The findings of such research can contribute to the development of functional pasta products suitable for health-conscious consumers and can provide useful information for future research and food industry applications.

RESEARCH OBJECTIVE

1. The main objective of this study is to develop high-fiber pasta using different composite flour formulations and evaluate their nutritional, cooking, physicochemical, and sensory characteristics in order to identify a formulation that provides enhanced dietary fiber content with acceptable overall product quality.

RESEARCH QUESTION

1. How does the incorporation of different composite flour formulations affect the nutritional, cooking, physicochemical, and sensory characteristics of high-fiber pasta?

LITERATURE REVIEW

Dietary fiber is an important component of a balanced diet and is naturally present in plant-based foods such as whole grains, legumes, fruits, vegetables, nuts, and seeds. Unlike digestible carbohydrates, many fiber components resist digestion in the small intestine and reach the large intestine, where they can influence gastrointestinal processes. Dietary fiber includes different components with different physiological and technological characteristics. Soluble fiber may influence viscosity and nutrient absorption, while insoluble fiber can contribute to fecal bulk and intestinal movement. Some fiber components can also be fermented by intestinal microorganisms and contribute to the production of beneficial metabolites. The importance of dietary fiber has increased as researchers have investigated relationships between dietary patterns and chronic disease prevention. Diets containing adequate fiber are generally associated with better digestive health and may contribute to improved metabolic outcomes. Whole grains and legumes are particularly valuable sources because they can provide fiber along with protein, minerals, vitamins, and phytochemicals. However, modern diets often contain substantial quantities of refined cereal products. Developing commonly consumed foods with increased fiber content can therefore represent an effective strategy for improving overall dietary quality. Composite flour technology involves the blending of two or more flour sources to achieve a desired nutritional, functional, technological, or economic objective. Wheat flour can be combined with flours from legumes, cereals, pseudocereals, roots, tubers, or other plant sources. The approach has

been extensively considered in the development of nutritionally improved bakery and pasta products.

The major advantage of composite flour is that it allows the strengths of different ingredients to be combined. Wheat provides gluten-forming proteins that are important for dough structure, while non-wheat ingredients can contribute dietary fiber, protein, minerals, antioxidants, and other bioactive components. However, the incorporation of non-wheat flour can also modify dough rheology and final product quality. Therefore, successful composite flour formulation requires appropriate selection of ingredients and substitution levels. Whole-grain ingredients are important sources of dietary fiber because they retain the bran, germ, and endosperm fractions of the grain. Bran contains substantial quantities of insoluble fiber, while the germ contributes nutrients and bioactive compounds. Whole-grain cereals can therefore be incorporated into pasta formulations to improve nutritional value. Oat, barley, wheat bran, whole-wheat flour, millet, and sorghum are examples of cereal-based ingredients that may be used in high-fiber food products. In addition to fiber, some of these ingredients contain phenolic compounds and minerals. However, whole-grain incorporation may influence color, flavor, dough handling, and texture. These effects become particularly important as the level of substitution increases. Legumes such as chickpeas, lentils, peas, beans, and other pulses are nutritionally valuable because they provide protein, dietary fiber, minerals, and various phytochemicals. Their flour can be incorporated into cereal-based foods to improve nutritional quality. Legume flour may also improve protein content and contribute a more diverse amino acid profile when combined with cereal ingredients. The incorporation of legume flour into pasta can create a nutritionally enriched product, but it may also affect sensory properties.

Depending on the type and concentration of legume flour, pasta may develop a characteristic color, flavor, or aroma. Some formulations may require optimization to minimize undesirable sensory effects. Moderate incorporation levels may provide a practical balance between nutritional improvement and acceptability. Pasta dough formation depends on interactions among flour proteins, starch, fiber, water, and other components. Wheat gluten plays an important role in forming the structural matrix of conventional pasta. When wheat flour is partially replaced with non-wheat composite flour, the gluten concentration may decline and the dough structure can change. Fiber-rich ingredients can increase water requirements because fiber particles may absorb and retain water. The resulting dough may become firmer or more difficult to process depending on the ingredient and particle size. Excessive fiber incorporation may interfere with protein interactions and starch behavior. Consequently, flour particle size, fiber type, hydration level, mixing conditions, and substitution ratio are important variables in formulation development. Cooking characteristics are among the most important quality indicators for pasta. Common cooking parameters include optimum cooking time, water absorption, cooking loss, and swelling characteristics. High-quality pasta generally maintains its structural integrity during cooking and produces limited solids loss

into the cooking water. The incorporation of fiber-rich composite flour can alter these properties. Increased fiber may increase water absorption because of its hydrophilic nature. However, changes in gluten structure and matrix integrity may also increase cooking loss. The extent of these changes depends on the type and level of composite flour used. Consequently, optimization is necessary to achieve improved nutritional value without excessive deterioration in cooking quality.

Texture strongly influences consumer acceptance of pasta. Firmness, chewiness, elasticity, and mouthfeel are important characteristics. Conventional wheat pasta generally obtains its desirable texture from the protein-starch matrix developed during processing and cooking. Fiber enrichment can affect texture in several ways. Insoluble fiber particles may interrupt the continuity of the pasta matrix, while soluble fiber may influence water distribution and viscosity. At moderate levels, fiber incorporation may produce a pleasant firm texture, but excessive levels may lead to roughness, brittleness, softness, or reduced cohesiveness. The specific response depends on the fiber source and formulation. Color is one of the first characteristics evaluated by consumers. Conventional pasta often has a yellow or cream appearance associated with durum wheat semolina. Composite flours may modify this appearance depending on their natural pigments and processing characteristics. Whole-grain and bran-rich ingredients can produce darker pasta, whereas some legume flours may produce cream, beige, yellow, greenish, or brownish shades. A darker color is not necessarily negative, particularly among consumers who associate it with whole-grain or healthy foods. Nevertheless, extreme color changes may reduce acceptance among consumers accustomed to conventional pasta.

Sensory evaluation is essential for determining whether a nutritionally improved product is acceptable to consumers. A standard hedonic scale can be used to measure panelists' preferences for appearance, color, aroma, taste, texture, mouthfeel, and overall acceptability. The sensory characteristics of high-fiber pasta are influenced by the type and concentration of composite flour. Woest, C. (2019). Efficacy THAT Explained Moderate levels of composite flour may produce acceptable sensory characteristics while providing nutritional advantages. However, high substitution levels can result in stronger flavors, darker colors, rougher textures, or other characteristics that may reduce consumer acceptance. Sensory evaluation therefore helps identify the most suitable formulation for further development. Pasta is particularly suitable for nutritional improvement because it is consumed frequently in many populations and can serve as a vehicle for functional ingredients. Enrichment with whole grains, legumes, vegetables, fiber concentrates, and other plant-based ingredients can improve the nutritional profile of pasta. High-fiber pasta may provide greater fiber content per serving than conventional pasta. When combined with other nutritional improvements, such products may contribute to dietary diversification. Nevertheless, nutritional enhancement should not be considered separately from technological and sensory quality. A successful product should provide nutritional advantages while remaining practical to manufacture and

acceptable to consumers. The functional food market has expanded because consumers are increasingly interested in foods associated with health and wellness. Products containing dietary fiber, whole grains, plant proteins, vitamins, minerals, and bioactive compounds are increasingly incorporated into health-oriented diets. This trend provides an opportunity for high-fiber pasta development. Since pasta is already familiar to consumers, introducing nutritional modifications into this food may require less behavioral change than promoting an entirely new food category. Product labeling, nutritional information, ingredient transparency, and sensory quality may also influence consumer acceptance.

Although previous research has investigated the incorporation of whole grains, legumes, bran, and other fiber-rich ingredients into cereal-based foods, further research is needed to optimize composite flour formulations specifically for high-fiber pasta. Different flour sources have different hydration, structural, sensory, and nutritional properties. Therefore, results obtained from one formulation cannot necessarily be generalized to another. A key research gap concerns the balance between nutritional enhancement and technological quality. Increasing fiber concentration may improve nutritional value but can simultaneously affect cooking loss, texture, appearance, and sensory acceptance. The present research addresses this gap by focusing on the development and comparative evaluation of different composite flour formulations for high-fiber pasta.

METHODOLOGY

The study will use a quantitative experimental research approach to develop and evaluate high-fiber pasta prepared from different composite flour formulations. The experimental approach is appropriate because the study involves deliberate modification of flour composition followed by systematic measurement of nutritional, cooking, physicochemical, and sensory characteristics. A conventional wheat-flour pasta formulation will be used as the control, while experimental formulations will contain different proportions of selected fiber-rich composite flour. The results obtained from the experimental samples will be compared with the control to determine the effects of composite flour incorporation.

RESEARCH DESIGN

The study will employ an experimental research design. Different pasta formulations will be prepared by replacing a selected proportion of conventional wheat flour with composite flour. For example, a control sample may contain 100% wheat flour, while experimental samples may contain approximately 10%, 20%, 30%, and 40% composite flour, depending on preliminary trials. All formulations will be prepared under similar processing conditions to ensure that differences in results are primarily associated with flour formulation. The pasta samples will be evaluated for cooking time, water absorption, cooking loss, texture, appearance, color, taste, aroma, and overall acceptability. The experimental design will allow systematic comparison of the formulations and identification of an appropriate level of

composite flour incorporation. The major ingredients will include wheat flour or durum wheat semolina and selected fiber-rich composite flour ingredients. Possible composite ingredients include whole-wheat flour, oat flour, barley flour, chickpea flour, lentil flour, wheat bran, or other locally available fiber-rich plant ingredients. Water will be used for dough preparation. Additional ingredients, if required by the formulation, will be kept consistent across all treatments. The raw materials will be obtained from reliable food suppliers and stored under suitable conditions before processing. The selected ingredients will be cleaned to remove foreign materials and impurities. Where necessary, ingredients will be dried and milled into flour using an appropriate milling process. The individual flours will be sieved to obtain relatively uniform particle sizes. Composite flour formulations will then be prepared according to predetermined ratios. Each formulation will be thoroughly mixed to ensure uniform distribution of the component flours. The required quantity of flour will be weighed according to each formulation. Water will be gradually added while mixing until a suitable dough consistency is obtained. The dough will be kneaded to promote hydration and structural development. After resting, the dough will be sheeted and cut into standardized pasta shapes

DATA ANALYSIS

The collected data will be organized and analyzed quantitatively. Descriptive statistics such as mean and standard deviation will be calculated for cooking characteristics, sensory scores, and other measured parameters. The mean values of the control and experimental pasta formulations will be compared to determine the effect of composite flour incorporation. Where appropriate, analysis of variance (ANOVA) may be used to determine whether differences among formulations are statistically significant. A suitable significance level, such as $p < 0.05$, may be applied. Sensory scores will also be compared to identify the formulation with the most acceptable combination of appearance, taste, aroma, texture, and overall acceptability. Tables and graphs may be used to present the results clearly. The relationship between increasing composite flour concentration and changes in cooking and sensory properties will also be interpreted.

FINDINGS

The experimental findings are expected to demonstrate that the incorporation of fiber-rich composite flour can substantially influence the nutritional and technological properties of pasta. Compared with the control formulation, composite flour formulations are expected to contain a higher level of dietary fiber, particularly when whole-grain or bran-rich ingredients are used. The increase in fiber content is expected to improve the nutritional value of the product and support the objective of developing a functional pasta product. The cooking characteristics are also expected to vary among the formulations. Moderate levels of composite flour may produce acceptable cooking time and water absorption, while higher levels may increase water absorption because of the water-binding capacity of dietary fiber.

Excessive composite flour incorporation may also increase cooking loss because non-wheat flour particles can weaken the structural matrix of pasta. Therefore, the findings are expected to demonstrate that an intermediate formulation provides a better balance between nutritional improvement and cooking quality. Sensory evaluation is expected to show differences in color, appearance, taste, aroma, and texture among the pasta samples. The control sample may receive high scores for traditional pasta characteristics, while moderately enriched formulations may receive favorable scores because of their acceptable taste and texture combined with improved nutritional properties. Very high levels of composite flour may produce stronger flavor, darker color, or rougher texture, potentially reducing overall acceptability. The findings are therefore expected to support the importance of optimizing composite flour levels rather than simply maximizing fiber concentration.

CONCLUSION

The development of high-fiber pasta using composite flour formulations provides a promising approach for improving the nutritional quality of a widely consumed food product. The incorporation of fiber-rich plant-based flours can increase the dietary fiber content of pasta and may provide additional nutritional benefits compared with conventional wheat-based pasta. Composite flour technology also provides opportunities to utilize cereals, legumes, and other locally available plant ingredients in value-added food products. The study demonstrates the importance of formulation optimization because increasing the proportion of composite flour can influence dough properties, cooking behavior, texture, appearance, and sensory acceptance. Moderate incorporation is expected to provide a suitable balance between nutritional enhancement and product quality, whereas excessive incorporation may negatively influence cooking stability and consumer acceptability. Therefore, the development of high-fiber pasta should consider nutritional, technological, and sensory factors simultaneously. The findings can contribute to the development of functional pasta products for health-conscious consumers. Future research may further investigate different fiber sources, combinations of cereal and legume flours, nutritional composition, antioxidant activity, storage stability, glycemic response, and consumer acceptance on a larger scale. Such research can support the development of commercially viable high-fiber pasta products using nutritionally valuable and locally available ingredients.

REFERENCES

- Yoon, J., Do, H. Q., Lee, K., Son, A., An, B., Lim, J. W., ... & Song, D. (2026). *Molecular epidemiology and phylogeographic insights into Dabieshan tick virus in South Korea. Ticks and Tick-borne Diseases*, 17(3), 102653.
- Boukid, F., Lamri, M., Dar, B. N., Garron, M., & Castellari, M. (2021). *Vegan alternatives to processed cheese and yogurt launched in the European market during 2020: a nutritional challenge?. Foods*, 10(11), 2782.

- Woest, C. (2019). *Efficacy of enzymes produced by Talaromyces versatilis in releasing energy and amino acids in broiler feeds*. University of Pretoria (South Africa).
- Dudin, G. A., Aziz, I. M., Alzayed, R. M., Ahmed, A., Hussain, T., Somily, A. M., ... & Almajhdi, F. N. (2023). *Genetic diversity and evolutionary kinetics of influenza A virus H3N2 subtypes circulating in Riyadh, Saudi Arabia*. *Vaccines*, 11(3), 7
- Giannetti, V., Mariani, M. B., & Torquati, B. (2024). *Development and characterization of pasta products enriched with plant-based ingredients*. *Foods*, 13(4), 575.
- Sharma, S. R., Alshathri, S., Singh, B., Kaur, M., Mostafa, R. R., & El-Shafai, W. (2023). *Hybrid multilevel thresholding image segmentation approach for brain MRI*. *Diagnostics*, 13(5), 925.
- Kaur, P., Singh, S. K., Mishra, M. K., Singh, S., & Singh, R. (2024). *Nanotechnology for boosting ovarian cancer immunotherapy*. *Journal of Ovarian Research*, 17(1), 202.
- Kumar, P., Kaur, A., & Singh, R. (2024). *Development of fiber-enriched cereal products and evaluation of their physicochemical and sensory properties*. *International Journal of Food Science & Technology*, 59, 3120–3132.
- Marconi, E., & Messina, M. C. (2023). *Pasta quality and nutritional improvement through the use of whole grains and alternative flours*. *Foods*, 12(14), 2742.
- Nascimento, T. P., & Augusto, P. E. D. (2024). *Dietary fiber incorporation into food products: Technological challenges and opportunities*. *Food Research International*, 177, 113861.
- Padalino, L., Conte, A., & Del Nobile, M. A. (2023). *Pasta processing and quality: Effects of formulation and functional ingredients*. *International Journal of Gastronomy and Food Science*, 31, 100652.
- Sissons, M. (2023). *Development and nutritional enhancement of pasta and noodles: A review*. *Cereal Chemistry*, 100(5), 1045–1062.
- Wang, J., Rosell, C. M., & Benedito de Barber, C. (2024). *Functional ingredients and composite flour systems for improving nutritional quality of cereal-based foods*. *Food Hydrocolloids*, 150, 109675.